

**TABLE 4**  
**Long-Term Trends in Lifetime Prevalence of Use of Various Drugs for Twelfth Graders**

	Percent ever used																									'00-'01 change			
	Class of 1975	Class of 1976	Class of 1977	Class of 1978	Class of 1979	Class of 1980	Class of 1981	Class of 1982	Class of 1983	Class of 1984	Class of 1985	Class of 1986	Class of 1987	Class of 1988	Class of 1989	Class of 1990	Class of 1991	Class of 1992	Class of 1993	Class of 1994	Class of 1995	Class of 1996	Class of 1997	Class of 1998	Class of 1999		Class of 2000	Class of 2001	
<i>Approx. N</i> <i>(in thousands) =</i>	9.4	15.4	17.1	17.8	15.5	15.9	17.5	17.7	16.3	15.9	16.0	15.2	16.3	16.3	16.7	15.2	15.0	15.8	16.3	15.4	15.4	14.3	15.4	15.2	13.6	12.8	12.8		
Any Illicit Drug <sup>a,b</sup>	55.2	58.3	61.6	64.1	65.1	65.4	65.6	64.4	62.9	61.6	60.6	57.6	56.6	53.9	50.9	47.9	44.1	40.7	42.9	45.6	48.4	50.8	54.3	54.1	54.7	54.0	53.9	-0.1	
Any Illicit Drug Other Than Marijuana <sup>a,b,c</sup>	36.2	35.4	35.8	36.5	37.4	38.7	42.8	41.1	40.4	40.3	39.7	37.7	35.8	32.5	31.4	29.4	26.9	25.1	26.7	27.6	28.1	28.5	30.0	29.4	29.4	29.0	28.5	-0.5	
Marijuana/Hashish	47.3	52.8	56.4	59.2	60.4	60.3	59.5	58.7	57.0	54.9	54.2	50.9	50.2	47.2	43.7	40.7	36.7	32.6	35.3	38.2	41.7	44.9	49.6	49.1	49.7	48.8	49.0	+0.2	
Inhalants <sup>d</sup>	—	10.3	11.1	12.0	12.7	11.9	12.3	12.8	13.6	14.4	15.4	15.9	17.0	16.7	17.6	18.0	17.6	16.6	17.4	17.7	17.4	16.6	16.1	15.2	15.4	14.2	13.0	-1.2	
Inhalants, Adjusted <sup>d,e</sup> Amyl/Butyl Nitrites <sup>f,g</sup>	—	—	—	—	18.2	17.3	17.2	17.7	18.2	18.0	18.1	20.1	18.6	17.5	18.6	18.5	18.0	17.0	17.7	18.3	17.8	17.5	16.9	16.5	16.0	14.6	13.8	-0.8	
Hallucinogens <sup>c</sup>	16.3	15.1	13.9	14.3	14.1	13.3	13.3	12.5	11.9	10.7	10.3	9.7	10.3	8.9	9.4	9.4	9.6	9.2	10.9	11.4	12.7	14.0	15.1	14.1	13.7	13.0	12.8	-0.2	
Hallucinogens, Adjusted <sup>h</sup> LSD	11.3	11.0	9.8	9.7	9.5	9.3	9.8	9.6	8.9	8.0	7.5	7.2	8.4	7.7	8.3	8.7	8.8	8.6	10.3	10.5	11.7	12.6	13.6	12.6	12.2	11.1	10.9	-0.2	
Hallucinogens Other Than LSD <sup>c</sup>	14.1	12.1	11.2	11.6	10.7	9.8	9.1	8.0	7.3	6.6	6.5	5.7	5.4	4.1	4.3	4.1	3.7	3.3	3.9	4.9	5.4	6.8	7.5	7.1	6.7	6.9	7.0	+0.1	
PCP <sup>g</sup>	—	—	—	—	12.8	9.6	7.8	6.0	5.6	5.0	4.9	4.8	3.0	2.9	3.9	2.8	2.9	2.4	2.9	2.8	2.7	4.0	3.9	3.9	3.4	3.4	3.5	+0.2	
MDMA (Ecstasy) <sup>f</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.1	6.9	5.8	8.0	11.0	11.7	+0.7
Cocaine	9.0	9.7	10.8	12.9	15.4	15.7	16.5	16.0	16.2	16.1	17.3	16.9	15.2	12.1	10.3	9.4	7.8	6.1	6.1	5.9	6.0	7.1	8.7	9.3	9.8	8.6	8.2	-0.4	
Crack <sup>i</sup>	—	—	—	—	—	—	—	—	—	—	—	—	5.4	4.8	4.7	3.5	3.1	2.6	2.6	3.0	3.0	3.3	3.9	4.4	4.6	3.9	3.7	-0.2	
Other Cocaine <sup>i</sup>	—	—	—	—	—	—	—	—	—	—	—	—	14.0	12.1	8.5	8.6	7.0	5.3	5.4	5.2	5.1	6.4	8.2	8.4	8.8	7.7	7.4	-0.3	
Heroin <sup>k</sup>	2.2	1.8	1.8	1.6	1.1	1.1	1.2	1.2	1.3	1.2	1.1	1.2	1.1	1.3	1.3	0.9	1.2	1.1	1.2	1.6	1.8	2.1	2.0	2.0	2.4	1.8	-0.6s		
With a needle <sup>l</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.7	0.8	0.9	0.8	0.9	0.8	0.7	-0.1	
Without a needle <sup>l</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.4	1.7	2.1	1.6	1.8	2.4	1.5	-0.9ss	
Other Narcotics <sup>m</sup>	9.0	9.6	10.3	9.9	10.1	9.8	10.1	9.6	9.4	9.7	10.2	9.0	9.2	8.6	8.3	8.3	6.6	6.1	6.4	6.6	7.2	8.2	9.7	9.8	10.2	10.6	9.9	-0.8	
Amphetamines <sup>b,m</sup>	22.3	22.6	23.0	22.9	24.2	26.4	32.2	27.9	26.9	27.9	26.2	23.4	21.6	19.8	19.1	17.5	15.4	13.9	15.1	15.7	15.3	15.3	16.5	16.4	16.3	15.6	16.2	+0.6	
Methamphetamine <sup>n</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Crystal Meth. (Ice) <sup>n</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.7	3.3	2.9	3.1	3.4	3.9	4.4	4.4	5.3	4.8	4.0	4.1	+0.1
Sedatives <sup>m,o</sup>	18.2	17.7	17.4	16.0	14.6	14.9	16.0	15.2	14.4	13.3	11.8	10.4	8.7	7.8	7.4	7.5	6.7	6.1	6.4	7.3	7.6	8.2	8.7	9.2	9.5	9.3	8.9	-0.4	
Barbiturates <sup>m</sup>	16.9	16.2	15.6	13.7	11.8	11.0	11.3	10.3	9.9	9.9	9.2	8.4	7.4	6.7	6.5	6.8	6.2	5.5	6.3	7.0	7.4	7.6	8.1	8.7	8.9	9.2	8.7	-0.5	
Methaqualone <sup>m,p</sup>	8.1	7.8	8.5	7.9	8.3	9.5	10.6	10.7	10.1	8.3	6.7	5.2	4.0	3.3	2.7	2.3	1.3	1.6	0.8	1.4	1.2	2.0	1.7	1.6	1.8	0.8	1.1	+0.3	
Tranquilizers <sup>c,m</sup>	17.0	16.8	18.0	17.0	16.3	15.2	14.7	14.0	13.3	12.4	11.9	10.9	10.9	9.4	7.6	7.2	7.2	6.0	6.4	6.6	7.1	7.2	7.8	8.5	9.3	8.9	9.2	+0.4	
Rohypnol <sup>f</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.2	1.8	3.0	2.0	1.5	1.7	+0.3
Alcohol <sup>q</sup>	90.4	91.9	92.5	93.1	93.0	93.2	92.6	92.8	92.6	92.6	92.2	91.3	92.2	92.0	90.7	89.5	88.0	87.5	87.0	—	—	—	—	—	—	—	—	—	—
Been Drunk <sup>n</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	65.4	63.4	62.5	62.9	63.2	61.8	64.2	62.4	62.3	62.3	63.9	+1.6
Cigarettes	73.6	75.4	75.7	75.3	74.0	71.0	71.0	70.1	70.6	69.7	68.8	67.6	67.2	66.4	65.7	64.4	63.1	61.8	61.9	62.0	64.2	63.5	65.4	65.3	64.6	62.5	61.0	-1.5	
Smokeless Tobacco <sup>r</sup>	—	—	—	—	—	—	—	—	—	—	—	31.4	32.2	30.4	29.2	—	—	32.4	31.0	30.7	30.9	29.8	25.3	26.2	23.4	23.1	19.7	-3.4	
Steroids <sup>n</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.0	2.9	2.1	2.1	2.0	2.4	2.3	1.9	2.4	2.7	2.9	2.5	3.7	+1.2s

NOTES: Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available.

Any apparent inconsistency between the change estimate and the prevalence of use estimates for the two most recent classes is due to rounding error.

SOURCE: The Monitoring the Future Study, the University of Michigan.