

TABLE 16
Long-Term Trends in Thirty-Day Prevalence of Use of Various Drugs for Twelfth Graders

	Percentage who used in last thirty days															
	Class of:															
Approx. N =	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	
Any Illicit Drug ^{a,b}	9400	15400	17100	17800	15500	15900	17500	17700	16300	15900	16000	15200	16300	16300	16700	
Any Illicit Drug Other Than Marijuana ^{a,b,c}	30.7	34.2	37.6	38.9	38.9	37.2	36.9	32.5	30.5	29.2	29.7	27.1	24.7	21.3	19.7	
Marijuana/Hashish	15.4	13.9	15.2	15.1	16.8	18.4	21.7	17.0	15.4	15.1	14.9	13.2	11.6	10.0	9.1	
Inhalants ^d	27.1	32.2	35.4	37.1	36.5	33.7	31.6	28.5	27.0	25.2	25.7	23.4	21.0	18.0	16.7	
Inhalants, Adjusted ^{d,e}	—	0.9	1.3	1.5	1.7	1.4	1.5	1.5	1.7	1.9	2.2	2.5	2.8	2.6	2.3	
Amyl/Butyl Nitrites ^{f,g}	—	—	—	—	3.2	2.7	2.5	2.5	2.5	2.6	3.0	3.2	3.5	3.0	2.7	
Hallucinogens ^c	—	—	—	—	2.4	1.8	1.4	1.1	1.4	1.4	1.6	1.3	1.3	0.6	0.6	
Hallucinogens, Adjusted ^{c,h}	4.7	3.4	4.1	3.9	4.0	3.7	3.7	3.4	2.8	2.6	2.5	2.5	2.5	2.2	2.2	
LSD	—	—	—	—	5.3	4.4	4.5	4.1	3.5	3.2	3.8	3.5	2.8	2.3	2.9	
Hallucinogens Other Than LSD ^c	2.3	1.9	2.1	2.1	2.4	2.3	2.5	2.4	1.9	1.5	1.6	1.7	1.8	1.8	1.8	
PCP ^{f,g}	3.7	2.3	3.0	2.7	2.4	2.3	2.1	1.7	1.5	1.6	1.3	1.3	1.1	0.7	0.8	
MDMA (Ecstasy) ^f	—	—	—	—	2.4	1.4	1.4	1.0	1.3	1.0	1.6	1.3	0.6	0.3	1.4	
Cocaine	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Crack ⁱ	1.9	2.0	2.9	3.9	5.7	5.2	5.8	5.0	4.9	5.8	6.7	6.2	4.3	3.4	2.8	
Other Cocaine ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	1.3	1.6	1.4	
Heroin ^k	—	—	—	—	—	—	—	—	—	—	—	—	4.1	3.2	1.9	
With a needle ^l	0.4	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.3	
Without a needle ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Other Narcotics ^{m,n}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Amphetamines ^{b,m}	2.1	2.0	2.8	2.1	2.4	2.4	2.1	1.8	1.8	1.8	2.3	2.0	1.8	1.6	1.6	
Methamphetamine ^o	8.5	7.7	8.8	8.7	9.9	12.1	15.8†	10.7	8.9	8.3	6.8	5.5	5.2	4.6	4.2	
Crystal Meth. (Ice) ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Sedatives (Barbiturates) ^m	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Sedatives, Adjusted ^{m,p}	4.7	3.9	4.3	3.2	3.2	2.9	2.6	2.0	2.1	1.7	2.0	1.8	1.4	1.2	1.4	
Methaqualone ^{m,q}	5.4	4.5	5.1	4.2	4.4	4.8	4.6	3.4	3.0	2.3	2.4	2.2	1.7	1.4	1.6	
Tranquillizers ^{c,m}	2.1	1.6	2.3	1.9	2.3	3.3	3.1	2.4	1.8	1.1	1.0	0.8	0.6	0.5	0.6	
Rohypnol ^f	4.1	4.0	4.6	3.4	3.7	3.1	2.7	2.4	2.5	2.1	2.1	2.1	2.0	1.5	1.3	
Alcohol ^f	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Been Drunk ^o	68.2	68.3	71.2	72.1	71.8	72.0	70.7	69.7	69.4	67.2	65.9	65.3	66.4	63.9	60.0	
Cigarettes	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Smokeless Tobacco ^{f,s}	36.7	38.8	38.4	36.7	34.4	30.5	29.4	30.0	30.3	29.3	30.1	29.6	29.4	28.7	28.6	
Steroids ^o	—	—	—	—	—	—	—	—	—	—	—	—	11.5	11.3	10.3	8.4
	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	

Cont'd

NOTE: See Table 14 for relevant footnotes.

TABLE 16 (cont'd)
Long-Term Trends in Thirty-Day Prevalence of Use of Various Drugs for Twelfth Graders

	Percentage who used in last thirty days															'03-'04 change
	Class of:															
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
Approx. N =	15200	15000	15800	16300	15400	15400	14300	15400	15200	13600	12800	12800	12900	14600	14600	
Any Illicit Drug ^{a,b}	17.2	16.4	14.4	18.3	21.9	23.8	24.6	26.2	25.6	25.9	24.9	25.7	25.4	24.1	23.4	-0.8
Any Illicit Drug Other Than Marijuana ^{a,b,c}	8.0	7.1	6.3	7.9	8.8	10.0	9.5	10.7	10.7	10.4	10.4†	11.0	11.3	10.4	10.8	+0.4
Marijuana/Hashish	14.0	13.8	11.9	15.5	19.0	21.2	21.9	23.7	22.8	23.1	21.6	22.4	21.5	21.2	19.9	-1.3
Inhalants ^d	2.7	2.4	2.3	2.5	2.7	3.2	2.5	2.5	2.3	2.0	2.2	1.7	1.5	1.5	1.5	0.0
Inhalants, Adjusted ^{d,e}	2.9	2.6	2.5	2.8	2.9	3.5	2.9	2.9	3.1	2.4	2.4	2.1	1.8	2.3	1.9	-0.4
Amyl/Butyl Nitrites ^{f,g}	0.6	0.4	0.3	0.6	0.4	0.4	0.7	0.7	1.0	0.4	0.3	0.5	0.6	0.7	0.7	0.0
Hallucinogens ^c	2.2	2.2	2.1	2.7	3.1	4.4	3.5	3.9	3.8	3.5	2.6†	3.3	2.3	1.8	1.9	+0.1
Hallucinogens, Adjusted ^{c,h}	2.3	2.4	2.3	3.3	3.2	4.6	3.8	4.1	4.1	3.9	3.0†	3.5	2.7	2.7	2.2	-0.5
LSD	1.9	1.9	2.0	2.4	2.6	4.0	2.5	3.1	3.2	2.7	1.6	2.3	0.7	0.6	0.7	+0.1
Hallucinogens Other Than LSD ^c	0.8	0.7	0.5	0.8	1.2	1.3	1.6	1.7	1.6	1.6	1.7†	1.9	2.0	1.5	1.7	+0.2
PCP ^{f,g}	0.4	0.5	0.6	1.0	0.7	0.6	1.3	0.7	1.0	0.8	0.9	0.5	0.4	0.6	0.4	-0.2
MDMA (Ecstasy) ^f	—	—	—	—	—	—	2.0	1.6	1.5	2.5	3.6	2.8	2.4	1.3	1.2	0.0
Cocaine	1.9	1.4	1.3	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.1	2.1	2.3	2.1	2.3	+0.3
Crack ⁱ	0.7	0.7	0.6	0.7	0.8	1.0	1.0	0.9	1.0	1.1	1.0	1.1	1.2	0.9	1.0	+0.1
Other Cocaine ⁱ	1.7	1.2	1.0	1.2	1.3	1.3	1.6	2.0	2.0	2.5	1.7	1.8	1.9	1.8	2.2	+0.4
Heroin ^k	0.2	0.2	0.3	0.2	0.3	0.6	0.5	0.5	0.5	0.5	0.7	0.4	0.5	0.4	0.5	+0.1
With a needle ^l	—	—	—	—	—	0.3	0.4	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	-0.1
Without a needle ^l	—	—	—	—	—	0.6	0.4	0.6	0.4	0.4	0.7	0.3	0.5	0.4	0.3	-0.1
Other Narcotics ^{m,n}	1.5	1.1	1.2	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.9	3.0†	4.0	4.1	4.3	+0.2
Amphetamines ^{b,m}	3.7	3.2	2.8	3.7	4.0	4.0	4.1	4.8	4.6	4.5	5.0	5.6	5.5	5.0	4.6	-0.3
Methamphetamine ^o	—	—	—	—	—	—	—	—	—	1.7	1.9	1.5	1.7	1.7	1.4	-0.3
Crystal Meth. (Ice) ^o	0.6	0.6	0.5	0.6	0.7	1.1	1.1	0.8	1.2	0.8	1.0	1.1	1.2	0.8	0.8	+0.1
Sedatives (Barbiturates) ^m	1.3	1.4	1.1	1.3	1.7	2.2	2.1	2.1	2.6	2.6	3.0	2.8	3.2	2.9	2.9	0.0
Sedatives, Adjusted ^{m,p}	1.4	1.5	1.2	1.3	1.8	2.3	2.3	2.1	2.8	2.8	3.1	3.0	3.4	3.0	2.9	-0.1
Methaqualone ^{m,q}	0.2	0.2	0.4	0.1	0.4	0.4	0.6	0.3	0.6	0.4	0.2	0.5	0.3	0.4	0.5	+0.1
Tranquilizers ^{c,m}	1.2	1.4	1.0	1.2	1.4	1.8	2.0	1.8	2.4	2.5	2.6†	2.9	3.3	2.8	3.1	+0.4
Rohypnol ^f	—	—	—	—	—	—	0.5	0.3	0.3	0.3	0.4	0.3	—	—	—	—
Alcohol ^f	57.1	54.0	51.3†	48.6	50.1	51.3	50.8	52.7	52.0	51.0	50.0	49.8	48.6	47.5	48.0	+0.5
Been Drunk ^o	—	31.6	29.9	28.9	30.8	33.2	31.3	34.2	32.9	32.9	32.3	32.7	30.3	30.9	32.5	+1.6
Cigarettes	29.4	28.3	27.8	29.9	31.2	33.5	34.0	36.5	35.1	34.6	31.4	29.5	26.7	24.4	25.0	+0.6
Smokeless Tobacco ^{f,s}	—	—	11.4	10.7	11.1	12.2	9.8	9.7	8.8	8.4	7.6	7.8	6.5	6.7	6.7	0.0
Steroids ^o	1.0	0.8	0.6	0.7	0.9	0.7	0.7	1.0	1.1	0.9	0.8	1.3	1.4	1.3	1.6	+0.3

SOURCE: The Monitoring the Future Study, the University of Michigan.