

TABLE 10
Smokeless Tobacco: Trends in 30-Day Prevalence of Use by Subgroups in Grade 10

	Percentage who used in last 30 days																					2011– 2012 change	
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	
<i>Approximate weighted N =</i>	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	16,200	16,100	15,100	15,900	15,200	14,900	15,000	
Total	10.0	9.6	10.4	10.5	9.7	8.6	8.9	7.5	6.5	6.1	6.9	6.1	5.3	4.9	5.6	5.7	6.1	5.0	6.5	7.5	6.6	6.4	-0.2
Gender:																							
Male	18.7	18.1	19.3	19.2	17.2	15.0	14.9	13.8	12.2	11.4	12.7	9.9	9.6	9.0	9.7	10.2	10.2	8.2	11.1	13.0	11.5	11.2	-0.3
Female	1.3	1.8	2.0	2.1	2.1	2.3	2.7	1.7	1.3	1.3	1.6	2.1	1.3	1.0	1.6	1.2	2.0	2.0	2.0	2.0	1.9	1.9	0.0
College Plans:																							
None or under 4 years	16.9	17.5	20.2	19.9	20.3	16.3	18.5	17.8	13.2	13.9	16.0	13.6	13.0	12.2	13.3	14.9	14.0	14.6	17.5	21.8	17.9	16.8	-1.2
Complete 4 years	8.4	8.0	8.4	8.5	7.8	7.2	7.2	5.7	5.4	4.8	5.4	4.8	4.1	3.9	4.5	4.5	5.0	3.8	5.3	5.8	5.2	5.3	+0.1
Region:																							
Northeast	8.6	5.3	8.0	9.0	7.6	6.8	9.3	6.5	5.2	4.6	4.9	4.7	4.5	5.1	4.6	4.5	5.9	3.3	6.6	6.7	5.6	4.3	-1.3
Midwest	11.0	9.6	10.0	10.0	11.0	9.5	7.1	7.9	8.1	6.2	7.0	4.8	4.9	3.7	5.7	7.5	6.3	7.2	7.7	9.2	6.4	7.9	+1.5
South	11.6	11.4	11.8	11.7	10.9	10.2	10.2	9.5	7.9	7.7	9.6	8.3	7.5	7.3	7.0	6.0	8.3	6.5	7.7	9.1	7.9	8.1	+0.2
West	7.8	10.9	11.1	10.9	7.7	6.0	8.2	4.6	4.0	4.5	3.0	5.1	3.5	3.0	4.5	4.1	3.3	2.2	3.3	4.0	5.4	4.0	-1.4
Population Density:																							
Large MSA	5.9	6.4	6.5	6.2	5.9	5.5	4.2	3.7	4.6	5.6	4.1	4.5	3.7	3.0	3.2	3.9	4.1	3.4	4.8	5.1	4.6	3.6	-1.1
Other MSA	9.2	9.3	10.1	10.9	9.2	8.4	8.3	5.7	5.3	4.3	5.7	6.1	4.8	4.3	5.3	5.3	5.8	4.4	6.3	7.0	6.1	6.3	+0.2
Non-MSA	14.7	13.3	14.1	13.9	15.0	12.2	14.7	15.1	11.3	9.8	12.5	8.2	9.2	9.0	9.4	9.6	9.9	8.7	9.3	12.1	10.2	10.5	+0.3
Parental Education: ^a																							
1.0–2.0 (Low)	6.6	10.1	10.9	9.4	9.6	8.1	9.0	6.8	7.2	7.4	6.9	6.7	7.4	3.7	4.4	7.4	6.1	5.0	7.9	6.5	6.4	7.2	+0.9
2.5–3.0	12.1	11.0	12.2	12.5	10.4	9.7	9.4	8.2	7.0	6.4	8.9	8.1	5.0	5.8	6.7	7.1	6.9	6.5	7.4	9.1	8.8	7.7	-1.1
3.5–4.0	10.6	10.5	10.9	10.2	10.9	8.3	10.3	8.6	7.3	6.3	7.1	5.5	4.9	5.2	6.0	5.7	6.4	5.4	7.4	8.7	7.2	7.5	+0.2
4.5–5.0	9.3	7.6	9.9	9.8	9.8	8.5	7.2	6.9	6.1	6.2	5.7	5.4	5.7	4.4	5.6	4.9	6.1	4.6	5.4	7.0	5.2	6.1	+0.9
5.5–6.0 (High)	8.6	8.1	7.0	8.9	6.0	7.7	8.3	5.2	4.8	4.0	4.8	5.2	4.3	4.4	3.8	5.3	4.8	2.9	5.6	6.0	5.3	3.8	-1.5
Race/Ethnicity (2-year average): ^b																							
White	—	11.4	12.0	12.5	12.0	11.0	10.4	10.0	8.7	7.5	7.5	7.8	6.9	6.1	6.6	6.9	7.3	7.3	7.5	9.1	9.1	8.0	-1.1
African American	—	2.9	2.3	2.3	2.5	2.5	2.8	2.3	1.6	2.0	3.2	2.6	2.5	2.7	2.5	3.2	3.3	2.8	2.6	2.3	2.2	2.3	0.0
Hispanic	—	6.2	6.1	4.3	3.6	4.0	4.6	4.8	4.8	4.5	4.0	4.0	4.1	3.3	3.1	3.4	2.7	2.6	2.9	2.8	3.4	4.4	+1.0

^aSource. The Monitoring the Future study, the University of Michigan.

^bNotes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See Table D-S2 in Appendix D of *Volume I* for the number of subgroup cases. See appendix B in *Volume I* for definition of variables in table. Data based on one of two forms in 1991–1996 and on two of four forms beginning in 1997; N is one half of N indicated in Table D-S2. In 2011 snus and dissolvable tobacco were added to the list of examples.

^aParental education is an average score of mother's education and father's education. See appendix B in *Volume I* for details.

^bTo derive percentages for each racial subgroup, data for the specified year and the previous year have been combined to increase subgroup sample sizes and thus provide more stable estimates. See appendix B in *Volume I* for details on how race/ethnicity is defined.