

TABLE 4
Trends in Prevalence of Use of *Smokeless Tobacco* in Grades 8, 10, and 12
 (Entries are percentages.)

	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	→ (Years cont.)
Lifetime																				
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	22.2	20.7	18.7	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	28.2	26.6	28.1	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	31.4	32.2	30.4	29.2	—	—	32.4	31.0	
30-Day																				
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.9	7.0	6.6	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	10.0	9.6	10.4	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	11.5	11.3	10.3	8.4	—	—	11.4	10.7	
Daily																				
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.6	1.8	1.5	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.3	3.0	3.3	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	4.7	5.1	4.3	3.3	—	—	4.3	3.3	
Approximate Ns																				
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	17,500	18,600	18,300	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	14,800	14,800	15,300	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	15,200	16,300	16,300	16,700	—	15,000	15,800	16,300	

(Table continued on next page.)

TABLE 4 (cont.)
Trends in Prevalence of Use of **Smokeless Tobacco in Grades 8, 10, and 12**
(Entries are percentages.)

	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2011– 2012 change
Lifetime																				
8th Grade	19.9	20.0	20.4	16.8	15.0	14.4	12.8	11.7	11.2	11.3	11.0	10.1	10.2	9.1	9.8	9.6	9.9	9.7	8.1	-1.6
10th Grade	29.2	27.6	27.4	26.3	22.7	20.4	19.1	19.5	16.9	14.6	13.8	14.5	15.0	15.1	12.2	15.2	16.8	15.6	15.4	-0.1
12th Grade	30.7	30.9	29.8	25.3	26.2	23.4	23.1	19.7	18.3	17.0	16.7	17.5	15.2	15.1	15.6	16.3	17.6	16.9	17.4	+0.5
30-Day																				
8th Grade	7.7	7.1	7.1	5.5	4.8	4.5	4.2	4.0	3.3	4.1	4.1	3.3	3.7	3.2	3.5	3.7	4.1	3.5	2.8	-0.7
10th Grade	10.5	9.7	8.6	8.9	7.5	6.5	6.1	6.9	6.1	5.3	4.9	5.6	5.7	6.1	5.0	6.5	7.5	6.6	6.4	-0.2
12th Grade	11.1	12.2	9.8	9.7	8.8	8.4	7.6	7.8	6.5	6.7	6.7	7.6	6.1	6.6	6.5	8.4	8.5	8.3	7.9	-0.4
Daily																				
8th Grade	1.9	1.2	1.5	1.0	1.0	0.9	0.9	1.2	0.8	0.8	1.0	0.7	0.7	0.8	0.8	0.8	0.9	0.8	0.5	-0.3
10th Grade	3.0	2.7	2.2	2.2	2.2	1.5	1.9	2.2	1.7	1.8	1.6	1.9	1.7	1.6	1.4	1.9	2.5	1.7	2.0	+0.3
12th Grade	3.9	3.6	3.3	4.4	3.2	2.9	3.2	2.8	2.0	2.2	2.8	2.5	2.2	2.8	2.7	2.9	3.1	3.1	3.2	+0.1
Approximate Ns																				
8th Grade	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	16,500	16,100	15,700	15,000	15,300	16,000	15,100	
10th Grade	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	16,200	16,100	15,100	15,900	15,200	14,900	15,000	
12th Grade	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200	14,500	14,000	13,700	14,400	14,100	13,700	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. For 8th and 10th graders: Data based on one of two forms for 1991–1996 and on two of four forms beginning in 1997; N is one half of N indicated. For 12th graders: Data based on one form; N is one fifth of N indicated in 1986–1988 and one sixth of N indicated beginning in 1989. The prevalence of use of smokeless tobacco was not asked of 12th graders in 1990 and 1991. Prior to 1990 the prevalence-of-use question on smokeless tobacco was located near the end of one 12th-grade questionnaire form, whereas after 1991 the question was placed earlier and in a different form. This shift could explain the discontinuities between the corresponding data. For all grades: In 2011 snus and dissolvable tobacco were added to the list of examples. An examination of the data did not show any effect from the wording change.