

TABLE 15
Long-Term Trends in Lifetime Prevalence of Use of Various Drugs in Grade 12

	Percentage who ever used																		→	
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	(Years cont.)
<i>Approximate weighted N =</i>	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200	15,000	15,800	16,300	
Any Illicit Drug ^{a,b}	55.2	58.3	61.6	64.1	65.1	65.4	65.6	64.4	62.9	61.6	60.6	57.6	56.6	53.9	50.9	47.9	44.1	40.7	42.9	
Any Illicit Drug other than Marijuana ^{a,b,c}	36.2	35.4	35.8	36.5	37.4	38.7	42.8	41.1	40.4	40.3	39.7	37.7	35.8	32.5	31.4	29.4	26.9	25.1	26.7	
Marijuana/Hashish	47.3	52.8	56.4	59.2	60.4	60.3	59.5	58.7	57.0	54.9	54.2	50.9	50.2	47.2	43.7	40.7	36.7	32.6	35.3	
Inhalants ^d	—	10.3	11.1	12.0	12.7	11.9	12.3	12.8	13.6	14.4	15.4	15.9	17.0	16.7	17.6	18.0	17.6	16.6	17.4	
Inhalants, Adjusted ^{d,e}	—	—	—	—	18.2	17.3	17.2	17.7	18.2	18.0	18.1	20.1	18.6	17.5	18.6	18.5	18.0	17.0	17.7	
Amyl/Butyl Nitrites ^{f,g}	—	—	—	—	11.1	11.1	10.1	9.8	8.4	8.1	7.9	8.6	4.7	3.2	3.3	2.1	1.6	1.5	1.4	
Hallucinogens ^c	16.3	15.1	13.9	14.3	14.1	13.3	13.3	12.5	11.9	10.7	10.3	9.7	10.3	8.9	9.4	9.4	9.6	9.2	10.9	
Hallucinogens, Adjusted ^{c,h}	—	—	—	—	17.7	15.6	15.3	14.3	13.6	12.3	12.1	11.9	10.6	9.2	9.9	9.7	10.0	9.4	11.3	
LSD	11.3	11.0	9.8	9.7	9.5	9.3	9.8	9.6	8.9	8.0	7.5	7.2	8.4	7.7	8.3	8.7	8.8	8.6	10.3	
Hallucinogens other than LSD ^c	14.1	12.1	11.2	11.6	10.7	9.8	9.1	8.0	7.3	6.6	6.5	5.7	5.4	4.1	4.3	4.1	3.7	3.3	3.9	
PCP ^{f,g}	—	—	—	—	12.8	9.6	7.8	6.0	5.6	5.0	4.9	4.8	3.0	2.9	3.9	2.8	2.9	2.4	2.9	
Ecstasy (MDMA) ^f	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Cocaine	9.0	9.7	10.8	12.9	15.4	15.7	16.5	16.0	16.2	16.1	17.3	16.9	15.2	12.1	10.3	9.4	7.8	6.1	6.1	
Crack ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	5.4	4.8	4.7	3.5	3.1	2.6	2.6	
Other Cocaine ^j	—	—	—	—	—	—	—	—	—	—	—	—	14.0	12.1	8.5	8.6	7.0	5.3	5.4	
Heroin ^k	2.2	1.8	1.8	1.6	1.1	1.1	1.1	1.2	1.2	1.3	1.2	1.1	1.2	1.1	1.3	1.3	0.9	1.2	1.1	
With a needle ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Without a needle ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Narcotics other than Heroin ^{m,n}	9.0	9.6	10.3	9.9	10.1	9.8	10.1	9.6	9.4	9.7	10.2	9.0	9.2	8.6	8.3	8.3	6.6	6.1	6.4	
Amphetamines ^{b,m}	22.3	22.6	23.0	22.9	24.2	26.4	32.2†	27.9	26.9	27.9	26.2	23.4	21.6	19.8	19.1	17.5	15.4	13.9	15.1	
Methamphetamine ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Crystal Methamphetamine (Ice) ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.7	3.3	2.9	3.1	
Sedatives (Barbiturates) ^{m,p}	16.9	16.2	15.6	13.7	11.8	11.0	11.3	10.3	9.9	9.9	9.2	8.4	7.4	6.7	6.5	6.8	6.2	5.5	6.3	
Sedatives, Adjusted ^{m,q}	18.2	17.7	17.4	16.0	14.6	14.9	16.0	15.2	14.4	13.3	11.8	10.4	8.7	7.8	7.4	7.5	6.7	6.1	6.4	
Methaqualone ^{m,r}	8.1	7.8	8.5	7.9	8.3	9.5	10.6	10.7	10.1	8.3	6.7	5.2	4.0	3.3	2.7	2.3	1.3	1.6	0.8	
Tranquilizers ^{c,m}	17.0	16.8	18.0	17.0	16.3	15.2	14.7	14.0	13.3	12.4	11.9	10.9	10.9	9.4	7.6	7.2	7.2	6.0	6.4	
Rohypnol ^f	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Alcohol ^s	90.4	91.9	92.5	93.1	93.0	93.2	92.6	92.8	92.6	92.6	92.2	91.3	92.2	92.0	90.7	89.5	88.0	87.5‡	80.0	
Been Drunk ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	65.4	63.4	62.5	
Cigarettes	73.6	75.4	75.7	75.3	74.0	71.0	71.0	70.1	70.6	69.7	68.8	67.6	67.2	66.4	65.7	64.4	63.1	61.8	61.9	
Smokeless Tobacco ^{ft}	—	—	—	—	—	—	—	—	—	—	—	31.4	32.2	30.4	29.2	—	—	32.4	31.0	
Steroids ^{m,u}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.0	2.9	2.1	2.1	2.0	

(Table continued on next page.)

TABLE 15 (cont.)
Long-Term Trends in Lifetime Prevalence of Use of Various Drugs in Grade 12

	Percentage who ever used																		2011– 2012 change	
	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	
<i>Approximate weighted N =</i>	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200	14,500	14,000	13,700	14,400	14,100	13,700	
Any Illicit Drug ^{a,b}	45.6	48.4	50.8	54.3	54.1	54.7	54.0	53.9	53.0	51.1	51.1	50.4	48.2	46.8	47.4	46.7	48.2	49.9	49.1	-0.8
Any Illicit Drug other than Marijuana ^{a,b,c}	27.6	28.1	28.5	30.0	29.4	29.4	29.0†	30.7	29.5	27.7	28.7	27.4	26.9	25.5	24.9	24.0	24.7	24.9	24.1	-0.9
Marijuana/Hashish	38.2	41.7	44.9	49.6	49.1	49.7	48.8	49.0	47.8	46.1	45.7	44.8	42.3	41.8	42.6	42.0	43.8	45.5	45.2	-0.3
Inhalants ^d	17.7	17.4	16.6	16.1	15.2	15.4	14.2	13.0	11.7	11.2	10.9	11.4	11.1	10.5	9.9	9.5	9.0	8.1	7.9	-0.2
Inhalants, Adjusted ^{d,e}	18.3	17.8	17.5	16.9	16.5	16.0	14.6	13.8	12.4	12.2	11.4	11.9	11.5	11.0	10.1	10.2	—	—	—	—
Amyl/Butyl Nitrites ^{f,g}	1.7	1.5	1.8	2.0	2.7	1.7	0.8	1.9	1.5	1.6	1.3	1.1	1.2	1.2	0.6	1.1	—	—	—	—
Hallucinogens ^c	11.4	12.7	14.0	15.1	14.1	13.7	13.0†	14.7	12.0	10.6	9.7	8.8	8.3	8.4	8.7	7.4	8.6	8.3	7.5	-0.8
Hallucinogens, Adjusted ^{c,h}	11.7	13.1	14.5	15.4	14.4	14.2	13.6†	15.3	12.8	10.9	9.9	9.3	8.8	8.9	9.0	8.0	9.1	8.8	7.9	-0.9
LSD	10.5	11.7	12.6	13.6	12.6	12.2	11.1	10.9	8.4	5.9	4.6	3.5	3.3	3.4	4.0	3.1	4.0	4.0	3.8	-0.2
Hallucinogens other than LSD ^c	4.9	5.4	6.8	7.5	7.1	6.7	6.9†	10.4	9.2	9.0	8.7	8.1	7.8	7.7	7.8	6.8	7.7	7.3	6.6	-0.8
PCP ^{f,g}	2.8	2.7	4.0	3.9	3.9	3.4	3.4	3.5	3.1	2.5	1.6	2.4	2.2	2.1	1.8	1.7	1.8	2.3	1.6	-0.7
Ecstasy (MDMA) ^f	—	—	6.1	6.9	5.8	8.0	11.0	11.7	10.5	8.3	7.5	5.4	6.5	6.5	6.2	6.5	7.3	8.0	7.2	-0.9
Cocaine	5.9	6.0	7.1	8.7	9.3	9.8	8.6	8.2	7.8	7.7	8.1	8.0	8.5	7.8	7.2	6.0	5.5	5.2	4.9	-0.3
Crack ⁱ	3.0	3.0	3.3	3.9	4.4	4.6	3.9	3.7	3.8	3.6	3.9	3.5	3.5	3.2	2.8	2.4	2.4	1.9	2.1	+0.2
Other Cocaine ^j	5.2	5.1	6.4	8.2	8.4	8.8	7.7	7.4	7.0	6.7	7.3	7.1	7.9	6.8	6.5	5.3	5.1	4.9	4.4	-0.4
Heroin ^k	1.2	1.6	1.8	2.1	2.0	2.0	2.4	1.8	1.7	1.5	1.5	1.5	1.4	1.5	1.3	1.2	1.6	1.4	1.1	-0.3
With a needle ^l	—	0.7	0.8	0.9	0.8	0.9	0.8	0.7	0.8	0.7	0.7	0.9	0.8	0.7	0.7	0.6	1.1	0.9	0.7	-0.2
Without a needle ^l	—	1.4	1.7	2.1	1.6	1.8	2.4	1.5	1.6	1.8	1.4	1.3	1.1	1.4	1.1	0.9	1.4	1.3	0.8	-0.5 s
Narcotics other than Heroin ^{m,n}	6.6	7.2	8.2	9.7	9.8	10.2	10.6	9.9†	13.5	13.2	13.5	12.8	13.4	13.1	13.2	13.2	13.0	13.0	12.2	-0.8
Amphetamines ^{b,m}	15.7	15.3	15.3	16.5	16.4	16.3	15.6	16.2	16.8	14.4	15.0	13.1	12.4	11.4	10.5	9.9	11.1	12.2	12.0	-0.2
Methamphetamine ^o	—	—	—	—	—	8.2	7.9	6.9	6.7	6.2	6.2	4.5	4.4	3.0	2.8	2.4	2.3	2.1	1.7	-0.3
Crystal Methamphetamine (Ice) ^o	3.4	3.9	4.4	4.4	5.3	4.8	4.0	4.1	4.7	3.9	4.0	4.0	3.4	3.4	2.8	2.1	1.8	2.1	1.7	-0.4
Sedatives (Barbiturates) ^{m,p}	7.0	7.4	7.6	8.1	8.7	8.9	9.2	8.7	9.5	8.8	9.9	10.5	10.2	9.3	8.5	8.2	7.5	7.0	6.9	-0.1
Sedatives, Adjusted ^{m,q}	7.3	7.6	8.2	8.7	9.2	9.5	9.3	8.9	10.2	9.1	10.1	11.0	10.6	9.6	8.9	8.4	7.6	7.2	7.2	+0.1
Methaqualone ^{m,r}	1.4	1.2	2.0	1.7	1.6	1.8	0.8	1.1	1.5	1.0	1.3	1.3	1.2	1.0	0.8	0.7	0.4	0.6	0.8	+0.2
Tranquilizers ^{c,m}	6.6	7.1	7.2	7.8	8.5	9.3	8.9†	10.3	11.4	10.2	10.6	9.9	10.3	9.5	8.9	9.3	8.5	8.7	8.5	-0.2
Rohypnol ^f	—	—	1.2	1.8	3.0	2.0	1.5	1.7	—	—	—	—	—	—	—	—	—	—	—	—
Alcohol ^s	80.4	80.7	79.2	81.7	81.4	80.0	80.3	79.7	78.4	76.6	76.8	75.1	72.7	72.2	71.9	72.3	71.0	70.0	69.4	-0.6
Been Drunk ^o	62.9	63.2	61.8	64.2	62.4	62.3	62.3	63.9	61.6	58.1	60.3	57.5	56.4	55.1	54.7	56.5	54.1	51.0	54.2	+3.2
Cigarettes	62.0	64.2	63.5	65.4	65.3	64.6	62.5	61.0	57.2	53.7	52.8	50.0	47.1	46.2	44.7	43.6	42.2	40.0	39.5	-0.5
Smokeless Tobacco ^{f,t}	30.7	30.9	29.8	25.3	26.2	23.4	23.1	19.7	18.3	17.0	16.7	17.5	15.2	15.1	15.6	16.3	17.6	16.9	17.4	+0.5
Steroids ^{m,u}	2.4	2.3	1.9	2.4	2.7	2.9	2.5	3.7	4.0	3.5	3.4	2.6	2.7	2.2	2.2	2.2	2.0	1.8	1.8	0.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. See footnotes following Table 18.