

TABLE 1
Annual Prevalence of Use for Various Types of Illicit Drugs, 2015
Among Full-Time College Students
1 to 4 Years beyond High School
by Gender

(Entries are percentages.)

	<u>Total</u>	<u>Males</u>	<u>Females</u>
Any Illicit Drug ^a	41.4	45.2	39.2
Marijuana	37.9	40.2	36.6
Any Illicit Drug other than Marijuana ^a	18.5	24.6	14.9
Adderall ^{b,f}	10.7	14.8	8.1
Cocaine	4.3	6.4	3.0
Hallucinogens ^c	4.3	6.6	3.0
Tranquilizers ^f	4.3	5.1	3.8
Ecstasy (MDMA) ^d	4.2	5.0	3.7
Other Cocaine ^d	4.2	6.4	3.0
Narcotics other than Heroin ^f	3.3	5.6	1.9
Hallucinogens other than LSD ^c	3.0	4.8	2.0
LSD ^c	3.0	4.9	2.0
Sedatives (Barbiturates) ^f	2.3	2.4	2.2
Ritalin ^{b,f}	2.0	3.2	1.2
Vicodin ^{b,f}	1.6	2.2	1.3
OxyContin ^{b,f}	1.5	3.2	0.4
Synthetic Marijuana ^b	1.5	1.6	1.5
Ketamine ^e	0.6	1.0	0.4
Methamphetamine ^e	0.5	1.3	*
Steroids ^e	0.3	0.7	*
Crack ^c	0.2	0.2	0.3
Bath Salts (synthetic stimulants) ^b	0.1	0.2	*
Heroin	0.1	0.3	0.1
GHB ^e	*	*	*
Crystal Methamphetamine (Ice) ^e	*	*	*
<i>Approximate Weighted N =</i>	<i>1,020</i>	<i>380</i>	<i>640</i>

Source. The Monitoring the Future study, the University of Michigan.

Notes. * * * indicates a prevalence rate of less than 0.05%.

See footnotes following Table 5.

TABLE 2
Annual Prevalence of Use for Alcohol and Tobacco, 2015
Among Full-Time College Students
1 to 4 Years beyond High School
by Gender

(Entries are percentages.)

	<u>Total</u>	<u>Males</u>	<u>Females</u>
Alcohol	79.0	78.7	79.2
Flavored Alcoholic Beverages ^h	64.5	50.3	72.7
Been Drunk ^b	61.6	64.1	60.0
Alcoholic Beverages containing Caffeine ^e	34.1	38.4	31.3
Tobacco using a Hookah ^b	23.4	24.5	22.8
Cigarettes	20.1	24.9	17.2
Small Cigars ^e	19.6	30.8	12.5
Snus ^e	5.8	14.8	*
Dissolvable Tobacco ^e	1.1	2.8	*
<i>Approximate Weighted N = 1,020</i>			

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' * ' indicates a prevalence rate of less than 0.05%.

See footnotes following Table 5.

TABLE 3
Thirty-Day Prevalence of Use for Various Types of Drugs, 2015
Among Full-Time College Students
1 to 4 Years beyond High School
by Gender

(Entries are percentages.)

	<u>Total</u>	<u>Males</u>	<u>Females</u>
Marijuana	21.1	25.5	18.5
Amphetamines, Adjusted ^{f,g}	4.2	6.6	2.7
Alcohol	63.2	63.9	62.7
Been Drunk ^b	38.4	44.4	34.4
Cigarettes	11.3	15.5	8.8
Electronic Vaporizers ^e	8.8	14.4	5.9
Large Cigars ^h	4.9	8.5	2.8
Flavored Little Cigars ^h	5.6	8.4	3.8
Regular Little Cigars ^h	4.1	7.2	2.2
<i>Approximate Weighted N =</i>	<i>1,020</i>	<i>380</i>	<i>640</i>

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' * ' indicates a prevalence rate of less than 0.05%.

See footnotes following Table 5.

TABLE 4
Thirty-Day Prevalence of Dailyⁱ Use for Various Types of Drugs, 2015
Among Full-Time College Students
1 to 4 Years beyond High School
by Gender

(Entries are percentages.)

	<u>Total</u>	<u>Males</u>	<u>Females</u>
Marijuana	4.6	6.7	3.4
Cigarettes	4.2	6.1	3.0
Alcohol	3.1	4.3	2.4
<i>Approximate Weighted N =</i>	<i>1,020</i>	<i>380</i>	<i>640</i>

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' * ' indicates a prevalence rate of less than 0.05%.

See footnotes following Table 5.

TABLE 5
Two-Week Prevalence of Binge Drinking, 2015
Among Full-Time College Students
1 to 4 Years beyond High School
by Gender

(Entries are percentages.)

	Total	Males	Females
5+ Drinks in a Row	31.9	37.3	28.6
<i>Approximate Weighted N =</i>	<i>1,020</i>	<i>380</i>	<i>640</i>
10+ Drinks in a Row*	11.2	18.3	6.9
15+ Drinks in a Row*	4.0	7.8	1.7
<i>Approximate Weighted N =</i>	<i>922</i>	<i>350</i>	<i>572</i>

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' * ' indicates a prevalence rate of less than 0.05%.

See footnotes on the following page.

*Due to limited sample size, numbers for 10+ drinks in a row and 15+ drinks in a row is based on data for years 2011-2015 combined.

Footnotes for Tables 1 through 5

^aUse of any illicit drug includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in three of the six questionnaire forms. Total *N* in 2015 for college students is approximately 510.

^cThis drug was asked about in five of the six questionnaire forms. Total *N* in 2015 for college students is approximately 850.

^dThis drug was asked about in four of the six questionnaire forms. Total *N* in 2015 for college students is approximately 680.

^eThis drug was asked about in two of the six questionnaire forms. Total *N* in 2015 for college students is approximately 340.

^fOnly drug use that was not under a doctor's orders is included here.

^gBased on the data from the revised question, which attempts to exclude inappropriate reporting of nonprescription amphetamines.

^hThis drug was asked about in one of the six questionnaire forms. Total *N* in 2015 for college students is approximately 170.

ⁱDaily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes, measured as actual daily use, and 5+ drinks, measured as having five or more drinks in a row in the last two weeks.