

**TABLE 1**  
**Annual Prevalence of Use for Various Types of Drugs, 2017:**  
**Full-Time College Students vs. Noncollege Youth**  
**among Respondents 1 to 4 Years beyond High School**  
**by Gender**

(Entries are percentages.)

	Total		Males		Females	
	Full-Time College	Non- College	Full-Time College	Non- College	Full-Time College	Non- College
Any Illicit Drug <sup>a</sup>	42.5	43.0	46.4	39.8	40.0	45.1
Any Illicit Drug other than Marijuana <sup>a</sup>	18.2	18.0	19.8	19.9	17.1	16.7
Marijuana	38.3	41.3	40.8	39.8	36.7	42.4
Synthetic Marijuana <sup>b</sup>	0.5	2.4	0.9	3.8	0.4	1.6
Inhalants <sup>b</sup>	1.7	0.6	2.3	1.5	1.4	*
Hallucinogens <sup>c</sup>	4.1	5.5	6.0	7.8	2.9	4.0
LSD <sup>c</sup>	2.9	4.0	4.1	5.7	2.1	2.9
Hallucinogens other than LSD <sup>c</sup>	2.5	3.2	3.3	5.3	2.0	1.9
MDMA (Ecstasy, Molly) <sup>d,e</sup>	2.6	4.7	3.1	4.2	2.2	5.1
Salvia <sup>b</sup>	0.3	2.5	0.9	1.4	*	3.2
Cocaine	4.8	5.2	6.6	3.9	3.6	6.0
Crack <sup>c</sup>	0.2	0.7	0.5	0.4	*	0.9
Other Cocaine <sup>d</sup>	4.5	4.8	5.2	3.9	4.0	5.3
Heroin	*	0.3	*	*	*	0.4
Narcotics other than Heroin <sup>f</sup>	3.1	4.1	3.7	4.2	2.7	4.1
OxyContin <sup>b,f</sup>	1.8	2.6	2.6	1.9	1.2	3.0
Vicodin <sup>b,f</sup>	1.1	1.8	1.4	2.1	0.9	1.6
Amphetamines, Adjusted <sup>f,g</sup>	8.6	7.3	7.7	7.5	9.2	7.2
Ritalin <sup>b,f</sup>	1.5	1.4	1.4	0.8	1.5	1.8
Adderall <sup>b,f</sup>	9.5	6.7	9.8	5.7	9.2	7.4
Bath Salts (synthetic stimulants) <sup>b</sup>	0.2	1.5	0.6	2.4	*	1.0
Sedatives (Barbiturates) <sup>f</sup>	1.9	2.4	2.4	3.1	1.6	1.9
Tranquilizers <sup>f</sup>	3.7	4.5	4.0	4.8	3.4	4.4
Alcohol	75.7	72.0	73.6	71.4	77.1	72.4
Been Drunk <sup>b</sup>	58.4	52.2	62.8	51.2	55.4	52.8
Cigarettes	16.6	30.4	20.0	35.5	14.4	26.9
Tobacco using a Hookah <sup>b</sup>	10.1	16.8	12.9	17.7	8.3	16.2
<i>Approximate Weighted N =</i>	873	498	344	201	529	296

Source. The Monitoring the Future study, the University of Michigan.

Notes. \* \* indicates a prevalence rate of less than 0.05%.

See footnotes following Table 3.

**TABLE 2**  
**Thirty-Day Prevalence of Use for Various Types of Drugs, 2017:**  
**Full-Time College Students vs. Noncollege Youth**  
**among Respondents 1 to 4 Years beyond High School**  
**by Gender**

(Entries are percentages.)

	Total		Males		Females	
	Full-Time College	Non- College	Full-Time College	Non- College	Full-Time College	Non- College
Any Illicit Drug <sup>a</sup>	23.5	27.2	26.4	27.7	21.5	27.0
Any Illicit Drug other than Marijuana <sup>a</sup>	7.0	8.4	7.2	9.4	6.9	7.8
Marijuana	21.3	27.6	24.4	28.8	19.3	26.8
Inhalants <sup>b</sup>	0.9	*	0.6	*	1.0	*
Hallucinogens <sup>c</sup>	1.2	1.5	1.2	2.1	1.3	1.1
LSD <sup>c</sup>	0.9	1.0	1.1	1.1	0.7	0.9
Hallucinogens other than LSD <sup>c</sup>	0.6	0.5	0.3	0.9	0.8	0.2
MDMA (Ecstasy, Molly) <sup>d,e</sup>	0.5	1.2	0.6	0.7	0.4	1.6
Cocaine	1.3	1.2	1.4	1.4	1.3	1.2
Crack <sup>c</sup>	*	0.4	*	*	*	0.7
Other Cocaine <sup>d</sup>	1.1	0.7	1.1	1.6	1.1	*
Heroin	*	0.3	*	*	*	0.4
Narcotics other than Heroin <sup>f</sup>	0.7	1.4	0.6	1.5	0.7	1.3
Amphetamines, Adjusted <sup>f,g</sup>	3.7	3.1	3.9	3.2	3.5	3.0
Sedatives (Barbiturates) <sup>f</sup>	0.5	0.9	0.7	1.2	0.4	0.8
Tranquilizers <sup>f</sup>	0.9	1.6	0.6	1.7	1.1	1.5
Alcohol	62.0	56.4	63.2	56.3	61.2	56.4
Been Drunk <sup>b</sup>	35.0	29.9	37.5	25.3	33.3	32.5
Cigarettes	7.9	21.7	11.3	24.4	5.8	19.9
<i>Approximate Weighted N =</i>	873	498	344	201	529	296

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' \* ' indicates a prevalence rate of less than 0.05%.

See footnotes following Table 3.

**TABLE 3**  
**Thirty-Day Prevalence of Daily<sup>h</sup> Use for Various Types of Drugs, 2017:**  
**Full-Time College Students vs. Noncollege Youth**  
**among Respondents 1 to 4 Years beyond High School**  
**by Gender**

(Entries are percentages.)

	Total		Males		Females	
	Full-Time College	Non- College	Full-Time College	Non- College	Full-Time College	Non- College
Marijuana	4.4	13.2	6.4	15.4	3.1	11.7
Alcohol						
Daily	2.2	4.0	3.7	8.2	1.1	1.1
5+ Drinks in a Row in Last 2 Weeks	32.9	28.1	38.6	31.5	29.2	25.8
Cigarettes						
Daily	2.0	14.4	3.4	14.5	1.1	14.4
1/2 Pack+/Day	0.2	7.9	0.6	7.2	*	8.3
<i>Approximate Weighted N =</i>	<i>873</i>	<i>498</i>	<i>344</i>	<i>201</i>	<i>529</i>	<i>296</i>

*Source.* The Monitoring the Future study, the University of Michigan.

*Notes.* \* \* indicates a prevalence rate of less than 0.05%.

See footnotes on the following page.

## Footnotes for Tables 1 through 3

<sup>a</sup>Use of any illicit drug includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

<sup>b</sup>This drug was asked about in three of the six questionnaire forms.

<sup>c</sup>This drug was asked about in five of the six questionnaire forms.

<sup>d</sup>This drug was asked about in four of the six questionnaire forms.

<sup>e</sup>Based on data from the revised question that includes "molly" as an example of MDMA

<sup>f</sup>Only drug use that was not under a doctor's orders is included here.

<sup>g</sup>Based on the data from the revised question, which attempts to exclude inappropriate reporting of nonprescription amphetamines.

<sup>h</sup>Daily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes, measured as actual daily use, and 5+ drinks, measured as having five or more drinks in a row in the last two weeks.